


1 / 3

F/G 20/11 NL


END
DATA
FILMED
7 83
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A129437

AFOSR-TR- 83 - 0497



Technical Report No. R-2-83

Project; Task No. 61102F; 2301/A3

Date: April 27, 1983

Title: Annual Report; Kinetic Theory

Contractor: Cornell University

Principal Investigator: Richard L. Liboff

DTIC
ELECTE
JUN 13 1983
A

Research supported in part by the Air Force Office of
Scientific Research under contract AFOSR-78-3574.

Distribution of this report is unlimited.

Approved for public release;
distribution unlimited.

83 06 10 147

DTIC FILE COPY

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER AFOSR-TR- 83 - 0497		2. GOVT ACCESSION NO. AD-A129437	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Annual Report			5. TYPE OF REPORT & PERIOD COVERED Annual [3/82 - 2/83]
7. AUTHOR(s) Richard L. Liboff			6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Cornell University Ithaca, N.Y. 14853			8. CONTRACT OR GRANT NUMBER(s) AFOSR 78-3574
11. CONTROLLING OFFICE NAME AND ADDRESS Directorate of Physics AFOSR Bolling AF Base, D.C. 20332			10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2301/A3
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)			12. REPORT DATE April 27, 1983
			13. NUMBER OF PAGES 9
			13. SECURITY CLASS. (of this report) Unclassified
			15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Approved for Public Release; Distribution Unlimited			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Short wavelength lasing; Population inversion; Dense reacting plasma; Coulomb gas; Matter-radiation interaction; Exciton-laser; Positronium laser; Collisional recombination.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A review of work performed under contract AFOSR 78-3574 during the '82-'83 support interval is presented. A list of titles and abstracts of technical reports issued during this period is included. A brief summary is presented of lectures delivered at the University of California on contractual research. The report concludes with a description of ongoing research.			

UNCLASSIFIED

Abstract

A review of work performed under contract AFOSR 78-3574 during the '82-'83 support interval is presented. A list of AF technical reports and invited talks on contractual research are included. The report concludes with a brief description of ongoing research.

Table of Contents

	<u>page</u>
Abstract	2
I. Technical Reports	3
II. Invited Talks	6
III. Ongoing Research	7

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
Distribution/	
Availability Codes	
Avail and/or	
Special	
A	

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TRANSMITTAL TO DTIC
 This technical report has been processed and is
 approved for public release in accordance with
 Distribution is unlimited.
MATTHEW J. KERPER
 Chief, Technical Information Division



I. Technical Reports

Titles and abstracts of technical reports issued during the '82-'83 support interval are listed below:

Report JQRST-1-82

"Review of Fundamental
Processes for Matter-Radiation
Interaction II"

January 28, 1982

Abstract

A concise review is presented of fundamental quantum electrodynamic processes which are relevant to X-ray lasers, gas lasers, and superpowerful lasers. Transition rates and cross sections are derived for atomic and free-electron Thomson scattering, Kramers-Heisenberg, and Raman scattering in the dipole approximation; Rayleigh scattering and Cerenkov effect. The report includes a table of reaction rates, cross sections, stopping power, and power spectra for all processes considered.

Report ZN-2-82

"Properties of a One-Dimensional
Coulomb Gas"

May 11, 1982

Abstract

The BBKGY equations for N identical, impenetrable, charged particles which move in one dimension and lie in a charge neutralizing background, are shown to separate into N uncoupled equations for the sequence of N reduced distributions. The potential relevant to any subgroup of s adjoining particles is that of an

s-dimensional harmonic oscillator whose frequency is the plasma frequency of the aggregate. The s-particle spatial equilibrium distribution reveals that particle vibrations remain centered about fixed, uniformly distributed sites as ρ/T goes from zero to infinity, where ρ is particle density and T is temperature. Thus it is concluded that the system suffers no change in phase for all ρ and T .

Report JPP-3-82

"Kinetic Theory for a Short-Wavelength
Lasing Plasma"

September 1, 1982

Abstract

A kinetic analysis is made of a reacting plasma dominated by three-body recombination and ionization, together with collisional and radiative excitation and de-excitation of atomic states. The plasma includes excited atoms, ions, electrons and photons. The kinetic theory yields rate equations for these species, together with explicit expressions for relevant rate coefficients. In the limit of spatial homogeneity and assuming atom and electron densities are close to equilibrium, an explicit form is obtained for the radiation absorption coefficient per unit length. A criterion is then constructed for population inversion. Application to a helium-like active medium (e.g. Al^{+11}) and hydrogen-like passive medium (e.g. Al^{+12}), at electron temperature of 300 eV, reveals that population inversion ensues at electron densities in excess of 10^{20} cm^{-3} . Algebraic solution of atomic state rate equations

demonstrates that the absorption coefficient grows insensitive to photon-atom interactions with increasing electron density.

Report R-4-82

"Exciton-Laser Amplifier"

December 1, 1982

Abstract

A laser-amplifying device is described which is based on the stimulated decay of excitons in a pure crystal. An estimate is made of the gain of the device. At a typical frequency the gain is found to be appreciably large thus suggesting practical application of the laser amplifier.

Report IJTP-1-83

"Induced Decay of Positronium
and Grasers"

April 22, 1983

Abstract

The differential cross section and the total cross section for the stimulated decay of positronium by an incident photon of frequency ω is calculated as a function of the dimensionless variable $\xi = \hbar\omega/mc^2$. For $\xi \gg 1$ the total cross section is found to decrease as ξ^{-2} . We also look at the particular case of positronium in a black-body radiation field. Expressions for the number of induced annihilations per second as functions of the dimensionless ratio mc^2/kT and the number of positronium atoms are obtained. It is found that this rate is proportional to $(kT/mc^2)^2$ for $kT \leq mc^2$ and to $(kT/mc^2)\ln(kT/mc^2)$ for $kT \gg mc^2$.

The possibility of utilizing induced two-photon decay of positronium as a γ -ray laser at the wavelength $\gamma_c/2$ is examined, where γ_c is the Compton wavelength.

II. Invited Talks

A lecture tour given by the principal investigator during August, 1982 proved to be particularly valuable to contractual research. Here is a brief summary of the highlights of this trip:

At UCLA, John Dawson discussed a short wavelength lasing scheme based on the metastable $2P-1S$ transition in H-like atoms. He further suggested that one of his earlier papers might prove relevant to the speaker's recent investigation of strongly coupled plasmas. At the end of the colloquium there was keen discussion on the proper choice of ionization coefficients for studies of a dense, recombining plasma. Neville Luhmann described present plasma work at UCLA.

At UCSD, Bill Thomson suggested that use of the Saha equation is somewhat inconsistent with population inversion studies since this equation implies a Boltzmann population of excited states. J. H. Malmberg described recent ongoing experiments in effecting a one-component plasma through use of a magnetic mirror device. It is believed that the plasma so confined will eventually reach the large γ domain. (Here γ denotes the plasma parameter.)

On Aug. 11 traveler met with Norman Rostoker (UC Irvine). Dr. Rostoker expressed great interest in recent strongly coupled

plasma analyses. He pointed out that research related to plasmas with $\gamma \approx 1$ would also prove valuable to the dynamics of the central region of the sun.

Attendance at all these talks was large and subsequent discussions were informative. There was valuable exchange of information with individual faculty members.

III. Ongoing Research

Present research is directed at the following areas:

1. Kinetic Theory of Strongly Coupled Plasmas.

In recombination lasing the plasma parameter, $\gamma \approx 1$, and the plasma is said to be strongly coupled. Kinetic properties of such plasmas permit the evaluation of parameters important to the design and implementation of short wavelength lasing.

2. Recombination Coefficient

Recombination and ionization coefficients are also relevant to recombination lasing. Save for J.J. Thomson's elementary result for recombination, formulations of this coefficient rest on gross assumptions concerning properties of a plasma. Present research in this area seeks to construct the recombination coefficient from first principles. This result would be particularly relevant to non-equilibrium plasmas.

3. Lasing Criterion

Research here seeks to obtain an explicit analytic expression for the integral criterion found by Heffernan and Liboff (Report No. JPP-3-82) for population inversion in a

dense recombining plasma. This newly constructed analytic form would easily permit graphical display which, in turn, would prove valuable to experimental research. Further study will generalize the Heffernan-Liboff criterion to include dynamics of the ion component of the plasma. Due to the complexity of this problem, influence of ion dynamics in the first formulation was minimal.